

more obscure and fine across from marginal vein, absent under submarginal. Hind wing dusky like fore wing, its marginal cilia as long as those of fore wing, its discal faint where present. Fore wing clear between stigmal vein and costal margin.

Little Mulgrave River, jungle, June 7, 1918.

Perilampoides tennysoni, new species.

Like *bicolor* but axillæ barely separated, thorax with no grooves; black purplish with head (save upper half of occiput and ocellar spots) and thorax save metathorax, meso-venter and meso-pleurum, a spot on each side of neck of pronotum and an oval spot on scutellum at meson between middle and apex, yellow brown; also antennæ, tibial tips, tibiæ 1, knees and tarsi. Funicle 6 abruptly enlarged to size of club, shorter than pedicel, others small, 1-5 slightly produced from one side at apex, somewhat wider than long, subequal, the ring-joint cup-shaped and nearly a half shorter. Truncate mandible 2 with its lateral apex acute. Stigmal long, equal marginal, post-marginal a fourth longer.

Two females taken from cavities in green twigs under green disc-galls placed side by side on their rims.

Gordonvale, June 16, 1918.

CORRECTIONS TO VOLUME VII

Page 155, line 31, for "tarsal" read *tarsals*; line 35, for "bend" read *band*.

Page 156, line 1, for "band" read *bands*; line 6, omit "only."

Page 157, line 12, for "penultimate" read *antepenultimate*.

Page 160, lines 8 and 9, for "dark with apical bands" read *with dark apical bands*.

C. S. LUDLOW.

CORRECTION OF PREOCCUPIED NAME

Mr. E. P. VanDuzee kindly calls my attention to the pre-occupation of *Grypotes* Dyar (Ins. Ins. Mens., v, 66, 1917) by *Grypotes* Fieber (Verh. zool.-bot. Ges. Wien, xvi, 503, 1866). The new name *Zatilpa* is accordingly suggested.

HARRISON G. DYAR.

Date of publication, February 27, 1920.